

# Final results of a Dark Matter Search with the Silicon Detectors of the CDMS II experiment and future results from SuperCDMS Soudan/SNOLAB

---

**Julien Billard**

Massachusetts Institute of Technology  
for the CDMS/SuperCDMS Collaborations

TAUP conference  
9th of September 2013



Massachusetts Institute of Technology

# The SuperCDMS Collaboration



California Inst. of Tech.  
B. Cornell, S.R. Golwala,  
D.C. Moore, R.H. Nelson



Fermi Nat. Accelerator Lab  
R. Basu Thakur D.A. Bauer,  
D. Holmgren, L. Hsu, B. Loer,



Massachusetts Inst. of Tech.  
A.J. Anderson, J. Billard,  
E. Figueroa-Feliciano, K.A. McCarthy



Pacific Northwest National  
Laboratory  
J. Hall



Queen's University  
C.H. Crewdson, P.C.F. Di Stefano,  
O. Kamaev, C. Martinez, P. Nadeau,  
K. Page, W. Rau, Y. Ricci



Santa Clara University  
B.A. Young



SLAC Nat. Accelerator Lab.  
M. Asai, A. Borgland,  
P.L. Brink, G.L. Godfrey,  
M.H. Kelsey, R. Partridge,  
K. Schneck, D.H. Wright



Southern Methodist University  
J. Cooley, B. Kara, H. Qiu, S. Scorza



Stanford University  
B. Cabrera, D.O. Caldwell\*,  
R.A. Moffat, P. Redl, B. Shank,  
S. Yellin, J.J. Yen



Syracuse University  
M.A. Bowles, R. Bunker, Y. Chen,  
M. Kiveni, R.W. Schnee



Texas A&M University  
H.R. Harris, A. Jastram,  
R. Mahapatra, K. Prasad



U. Autónoma de Madrid  
D. G. Cerdeno, L. Esteban,  
E. Lopez Asamar



University of British Columbia  
S.M. Oser, H. Tanaka



U. of California, Berkeley  
M. Daal, T. Doughty,  
N. Mirabolfathi, A. Phipps, M. Pyle,  
B. Sadoulet, B. Serfass, D. Speller



U. of Colorado, Denver  
M.E. Huber



University of Evansville  
A. Reisetter



University of Florida  
R. Agnese, D. Balakishiyeva,  
T. Saab, B. Welliver



University of Minnesota  
H. Chagani, P. Cushman, S. Fallows,  
T. Hofer, A. Kennedy,  
K. Koch, V. Mandic, M. Pepin,  
A.N. Villano, J. Zhang



University of South Dakota  
J. Sander



# The SuperCDMS Collaboration

Time 

*See R. Nelson talk*

## CDMS II (Ge+Si)

- 4.6 kg Ge (19 x 240 g)
- 1.2 kg Si (11 x 106g)
- 35% NR acceptance




## SuperCDMS Soudan

- Increased mass: 9.0 kg Ge (15 x 600 g)
- Increased acceptance
- Improved surface event discrimination

## SuperCDMS SNOLAB

- Proposed 200kg Ge array
  - Extensive R&D underway
  - Scale to 1 kg crystals
- Projected sensitivity of  $8 \times 10^{-47} \text{ cm}^2$

CDMSLite: See J. Hall talk




# CDMS II Silicon: Final analysis

See R. Nelson talk

## CDMS II (Ge+Si)

- 4.6 kg Ge (19 x 240 g)
- 1.2 kg Si (11 x 106g)
- 35% NR acceptance


Blind analysis of 140 kg-days of  
Si data (8 detectors)  
July 2007- September 2008



## SuperCDMS Soudan


- Increased mass: 9.0 kg Ge (15 x 600 g)
- Increased acceptance
- Improved surface event discrimination

CDMSLite: See J. Hall talk




## SuperCDMS SNOLAB

- Proposed 200kg Ge array
  - Extensive R&D underway
  - Scale to 1 kg crystals
- Projected sensitivity of  $8 \times 10^{-47} \text{ cm}^2$








# CDMSII ZIP detectors




240 g Ge or 106 g Si crystals  
(1 cm thick, 7.5 cm diameter)

- **Z**-sensitive **I**onization and **P**honon mediated
- Measure both **charge** and **phonon** energy
- Photolithographically patterned to collect **athermal** phonons and ionization signals
- Direct xy-position imaging
- Surface (z) event rejection from pulse shapes and timing



# CDMS strategy

$$Y = E_{\text{charge}}/E_{\text{recoil}}$$




Ionization yield (ionization energy per unit recoil energy) strongly depends on recoil type.

Most backgrounds produce electron recoils:  $Y = 1$

WIMPs and neutrons produce nuclear recoils:  $Y \sim 0.3$



Surface events: reduced ionization yield  $Y < 1$



Can be rejected through a phonon timing cut


Ionization yield + timing result in  $< 1$  in  $10^6$  electron recoils leaking into the signal region

pre-unblinding leakage estimate of 0.47






# Unblinding Results - before timing cut




# Unblinding Results - after timing cut





# Three events!



Shades of blue indicate the three separate timing cut energy ranges.



# Post-Unblinding Checks

- Consistency checks
  - Events occurred during high-quality data series
  - Events were well-reconstructed
  - Checked energy in other detectors to verify events were single scatters
- Updated background estimates:
  - Cosmogenic and radiogenic neutron:  
 **$<0.13$  @ 90% C.L.**
  - Surface events  


$$0.41^{+0.20}_{-0.08}(\text{stat.})^{+0.28}_{-0.24}(\text{syst.}) \longrightarrow$$
  - Possible Pb-206 recoils from Po-210 decays  
 **$<0.08$  @ 90% C.L.**





# Profile Likelihood analysis

- The maximum likelihood occurs at a WIMP mass of  $8.6 \text{ GeV}/c^2$  and WIMP-nucleon cross section of  $1.9 \times 10^{-41} \text{ cm}^2$
- Probability of observing 3 or more events from background fluctuations is equal to 5.4%
- Goodness of fit of the WIMP +Background model is 68.6%
- A profile likelihood ratio test statistic favors the WIMP +Background hypothesis over the background only at 99.81% C.L.



**We do not believe this result rises to the level of a discovery, but does call for further investigation.**

# SuperCDMS Soudan

See R. Nelson talk  
↙

## CDMS II (Ge+Si)

- 4.6 kg Ge (19 x 240 g)
- 1.2 kg Si (11 x 106g)
- 35% NR acceptance



## SuperCDMS Soudan

- Increased mass: 9.0 kg Ge (15 x 600 g)
- Increased acceptance
- Improved surface event discrimination

## SuperCDMS SNOLAB

- Proposed 200kg Ge array
  - Extensive R&D underway
  - Scale to 1 kg crystals
- Projected sensitivity of  $8 \times 10^{-47} \text{ cm}^2$



Installation complete Nov. 8, 2011. Detectors have been operating in DM-search mode since March 2012.






# SuperCDMS Soudan

- Array of 15 iZIPs in the Soudan infrastructure built for CDMS-II
- Factor  $> \times 10$  sensitivity increase over CDMS-II
  - Larger detector mass (x2.5 thicker detectors)
  - Fiducial fraction improved to  $\sim 50\%$  from 35%
  - Surface background negligible
- Interleaved technology allows to get:
  - Complex E-field to better tag surface events
  - 2 charge electrodes on each side (1 inner + 1 outer)
  - 4 phonon channels on each side (3 inner + 1 outer)



# SuperCDMS Soudan: charge discrimination


- Surface events (alpha, betas, nuclei) have **1-sided** charge collection
- Bulk events (gammas, nuclear recoils) have **2-sided** charge collection
- Charge **symmetry** is a powerful tag of surface events on top and bottom sides



# SuperCDMS Soudan: charge discrimination


- ~71,000 electrons and ~16,200  $^{206}\text{Pb}$  recoil surface events collected from  $^{210}\text{Pb}$  source.

- No events leaking into the signal region into ~50% fiducial volume (8-115 keVnr) in 37.6 live time days (March - July 2012)



- Limits surface events leakage to  $1.7 \times 10^{-5}$  @90% C.L. from 8 to 115 keVnr

- Good enough for a 0.3 ton-year exposure for SuperCDMS@ SNOLAB!



R. Agnese et al., arXiv:1305.2405

$^{206}\text{Pb}$  recoils

# SuperCDMS Soudan: phonon fiducialisation


Complex electric field creates asymmetry in both Ionization and **phonon** signals for surface events

*Surface event at **side 1**  
4.5 keV phonon energy*


- Surface event discrimination using phonon is much more significant at low energies
- Phonon baseline noise is lower than the charge one

**Promising for the Low Threshold analysis**  
*Not needed for the High Threshold analysis*

Charge traces



Side summed phonon traces





# SuperCDMS SNOLAB

See R. Nelson talk

## CDMS II (Ge+Si)



- 4.6 kg Ge (19 x 240 g)
- 1.2 kg Si (11 x 106g)
- 35% NR acceptance

## SuperCDMS Soudan

- Increased mass: 9.0 kg Ge (15 x 600 g)
- Increased acceptance
- Improved surface event discrimination


## SuperCDMS SNOLAB

- Proposed 200kg Ge array
  - Extensive R&D underway
  - Scale to 1 kg crystals
- Projected sensitivity of  $8 \times 10^{-47} \text{ cm}^2$




# SuperCDMS SNOLAB

- SuperCDMS has proposed a 200 kg Germanium advanced ***interleaved detector*** array for the SNOLAB facility
- Now, seriously considering adding some ***Si iZIPs***




First 100 mm iZIP has been fabricated with new phonon layout. Detector R&D underway



- With such an increase in exposure, we need to go deeper: ***from Soudan to SNOLab***.
- Reduction of cosmogenic neutron background by a factor of 100

# SuperCDMS SNOLAB



# Conclusion

---

## CDMSII final Si exposure

- Analysis of a 140.23 kg-day exposure of the CDMS-II Si detectors has been performed.
- Three events were seen in the signal region with a total expected background of 0.7 events.
- A profile likelihood analysis favors a WIMP+background hypothesis over the known background estimate as the source of our signal at the 99.81% confidence level ( $\sim 3\sigma$ , p-value: 0.19%).
- We do not believe this result rises to the level of a discovery, but does call for further investigation.

## SuperCDMS Soudan/SNOLAB

- SuperCDMS-Soudan ( $\sim 9$  kg) is taking data with iZIP detectors and expects to reach a WIMP-nucleon sensitivity of  $2 \times 10^{-45} \text{ cm}^2$  for spin-independent interactions.
- We have demonstrated surface event rejection with the new iZIP detector design using  $^{210}\text{Pb}$  sources down to  **$1.7 \times 10^{-5}$  @90% C.L.** which paves the way for below than  $10^{-46} \text{ cm}^2$  sensitivity at SNOLAB.
- Additional background rejection could be done using phonon pulse shape discrimination (study ongoing)